



Litacold[®]

BABY NASAL SPRAY

USAGE



Lay the baby down (or have the child sit up).
Turn the head to the side.



Gently insert the tip into the upper nostril and spray Litacold Baby and repeat the process in the other nostril.



Straighten the head to allow the mucus to drain.

With infants: wipe away any excess fluid. Use a baby nasal aspirator in the event of high flow.

With young children: ask them to blow their nose.



Wash the tip with hot soapy water after each use. Rinse and wipe.



RECOMMENDATION: 1 spray per nostril, preferably before meal.
1 to 4 times a day as necessary.

GENERIC HEALTHCARE PVT. LTD.
29A, Shivaji Nagar, Pune-411005.
INDIA.
export@ghpl.co
© Marque déposée / Trade mark.

16/0666



Litacold[®]

SPRAY NASAL BÉBÉ

USAGE



Allonger ou faire asseoir l'enfant
Incliner légèrement la tête sur le côté



Inserer doucement l'embout dans la narine supérieure et vaporiser Litacold Bébé et répéter le processus dans l'autre narine.



Redresser la tête pour permettre au mucus de s'écouler.

Avec les nourrissons : essuyer tout excès de liquide. Utiliser un aspirateur nasal pour bébé en cas de débit élevé.

Avec de jeunes enfants : leur demander de se moucher.



Laver l'embout à l'eau chaude savonneuse après chaque utilisation. Rincer et essuyer.



RECOMMANDATION :
1 pulvérisation par narine, de préférence avant les repas.
1 à 4 fois par jour selon les besoins.

GENERIC HEALTHCARE PVT. LTD.
29A, Shivaji Nagar, Pune-411005.
INDIA
export@ghpl.co
© Marque déposée / Trade mark.

16/0666